

## One-Sample Methods

PSY 5101: Advanced Statistics for  
Psychological and Behavioral Research I

## New Test Statistics

- All test statistics (inferential methods) have some things in common: use of descriptive statistics, use of probability...all of the basics of hypothesis testing. For example, all have a null hypothesis, all use  $\alpha$ , and for all, increasing N increases power
- But some things are different. For every new test statistic, we will cover four topics:
  - Situation, including the hypotheses
  - Test statistic
  - Theoretical reference distribution, critical values, and decision rules
  - Assumptions

## New Test Statistics

- I encourage you to start a chart. Put the four topics on the left side (rows) and the test statistics on the top (columns). Start with  $z_{\bar{X}}$ .

	$z_{\bar{X}}$
1. Situation/hypotheses	One sample $H_0: \mu = 100$ $\sigma^2$ known
2. Test statistic	$z_{\bar{X}} = \frac{\bar{X} - \mu}{\frac{\sigma}{\sqrt{N}}}$
3. Distribution	$N(0, 1)$
4. Assumptions	1. Population is normal 2. Observations are independent

## One-Sample t-Test

One-sample t	
1. Situation/hypotheses	One sample $H_0: \mu = 100$ $\sigma^2$ unknown
2. Test statistic	$t = \frac{\bar{X} - \mu}{\frac{s^2}{\sqrt{N}}}$
3. Distribution	$t_{df=N-1}$
4. Assumptions	1. Population is normal 2. Observations are independent Same as for $Z_{\bar{X}}$

---

---

---

---

---

---


---

---


## t Distributions

### • t distributions have the following characteristics:

- Theoretical distributions that are symmetric, smooth, unimodal, and have  $\mu=0$  (family of t distributions)
- Looks like the standard normal distribution but has longer tails and more variability
- The greater variability is due to t statistics having not only a mean ( $\bar{X}$ ) that varies from sample to sample but also a variance ( $s^2$ )



As df increases, the tails change such that the probability area in the tails decreases




---

---

---

---

---

---

---

---

## t Distributions: Degrees of Freedom

### • t distributions have only one parameter: degrees of freedom (df)

- Definition of df: parameter of a theoretical distribution
- The formula for df can change from one t statistic to the next
- df is directly tied to the amount of variability in the tail(s) of the distribution

### • The working definition for df is "in an estimate of variability, df is equal to the number of independent components minus the number of parameters estimated"

- To find the df for a test statistic, look at the estimate of variability that it uses and find the "independent components" and "number of parameters estimated"

### • The one-sample t has the unbiased sample variance ( $s^2$ ) in its formula

- In  $s^2 = \frac{\sum (X - \bar{X})^2}{N - 1}$ , there are N values of X (the independent components) and 1 statistic ( $\bar{X}$ ) that estimates the 1 parameter ( $\mu$ )
- So  $df = N - 1$  for the one-sample t

---

---

---

---

---

---

---

---

## Degrees of Freedom for One-Sample $t$

- The df for the one-sample  $t$  is  $N-1$
- Note that the whole concept of df came with the  $t$ -test
- There was no concept of df associated with  $z_{\bar{X}}$
- So whatever changed from  $z_{\bar{X}}$  to  $t$  is what brought with it the concept of df
- So how does  $t$  differ from  $z_{\bar{X}}$ ?

- $t$  has  $s^2$

$$z_{\bar{X}} = \frac{\bar{X} - \mu}{\frac{\sigma^2}{\sqrt{N}}}$$

No  $s^2$  and No df

$$t = \frac{\bar{X} - \mu}{\frac{s^2}{\sqrt{N}}}$$

$s^2$  and df


- The whole formula for degrees of freedom ( $df=N-1$ ) goes with  $s^2$
- Degrees of freedom are a way of "keeping score" by accounting for how many elements are allowed to vary

## One-Sample $t$ -Test: Example

- Are people who are interrupted in a task accurate in estimating how long they have spent on the task?
- People who were given 20 3-letter anagrams to solve (e.g., *arn* is *ran*) were interrupted after doing 10 of them and asked to estimate how long they had worked on the task
- The researchers formed a ratio of estimated to actual time, and  $\mu_{\text{ratio}}$  should be 1 if the people are accurate in estimating time
- The ratios for the  $N=10$  people are .911 1.011 1.807 2.010 1.911 2.156 1.251 1.516 2.730 1.160

## $t$ Table

- Now we can use df and  $\alpha=.05$  to find a critical value for  $t$  ( $t_{\text{crit}}$ )
- The  $t$  table is organized by df for the rows and  $\alpha$  for one- and two-tailed tests for the columns
- If  $N=10$ , then for a one-sample  $t$ ,  $df=N-1=10-1=9$
- For a two-tailed test with  $\alpha=.05$  and  $df=9$ , the critical values are  $\pm 2.262$



## One-Sample t-Test: Example

- Now compute the mean ( $\bar{X}$ ) and the unbiased variance ( $s^2$ )
  - $\bar{X} = 1.646$
  - $s^2 = .3352$
- Hypotheses:
  - $H_0: \mu = 1$
  - $H_1: \mu \neq 1$
- So we are now ready to compute  $t = \frac{\bar{X} - \mu}{\frac{s}{\sqrt{N}}} = \frac{1.646 - 1}{\sqrt{\frac{.3352}{10}}} = 3.53$
- Using a critical value decision rule, the upper  $t_{crit}$  is 2.262 and  $3.53 > 2.262$
- Using a p-value decision rule, the SPSS p-value was  $.0064 < \alpha = .05$
- So both decision rules lead us to reject  $H_0: \mu = 1$
- What does this look like in the sampling distribution of  $t$ ?

---

---

---

---

---

---


---

---

## One-Sample t-Test: Example

- Find the observed  $t = 3.53$  and the upper  $t_{crit} = 2.262$  in the distribution below
- Because  $3.53 > 2.262$  (or because  $.0064 < .05$ ) we reject  $H_0: \mu = 1$
- People interrupted in a task significantly overestimate the time spent in the task

Reporting style: "The ratio of estimated to actual time to solve the first 10 anagrams in the 20 anagram list was significantly greater than 1.0,  $\bar{X} = 1.646$ ,  $s = .579$ ,  $t(9) = 3.53$ ,  $p < .0064$ ."




---

---

---

---

---

---

---

---

## Test of Correlation: r

- Continuing with your chart, we will add a new test statistic to  $z_{\bar{X}}$  and the one-sample  $t$
- You already know it as a descriptive statistic but here it will be used to test hypotheses

1. Situation/hypotheses

2. Test statistic

3. Distribution

4. Assumptions

One sample

$H_0: \rho = 0$

X, Y pairs

$$r = \frac{\text{Cov}(X, Y)}{\sqrt{\text{Var } X} \sqrt{\text{Var } Y}}$$

$r_{df=N-2}$

1. Population is bivariate normal

2. Subjects are independent

---

---

---

---

---

---

---

---

## Test of Correlation: Degrees of Freedom

- The df for  $r$  is  $N-2$ 
  - It can be shown why  $df=N-2$  from the standard error of estimate
- The standard error of estimate is a statistic that describes spread of errors (or  $Y$  scores) in correlation and regression
- So, in  $s_{y.x}$  we look for independent components and statistics (that estimate parameters)

$$s_{y.x} = \sqrt{\frac{\sum(Y-Y')^2}{N-2}}$$

- The  $N$  values of  $Y$  are the independent components and  $Y'=bX+a$  has two statistics ( $b$  and  $a$ )
- So  $df=N-2$

---

---

---

---

---


---

---

---

## $r$ Critical Values

- Now we can use  $df$  and  $\alpha=.05$  to find a critical value for  $r$  ( $r_{crit}$ )
- The table of  $r_{crit}$  is organized by  $df$  for the rows and  $\alpha$  for one- and two-tailed tests for the columns
- If  $N=10$ , then for  $r$ ,  $df=N-2=10-2=8$
- For a two-tailed test with  $\alpha=.05$  and  $df=8$ , the critical values are  $\pm.632$




---

---

---

---

---

---

---


---

## Test of Correlation: Example

- Researchers believed stress for police officers is associated with the number of hours spent moonlighting on a second job
- For 28 officers,  $r$  was .45
- Is  $r$  significantly different from zero?

$H_0: \rho=0$ ,  $H_1: \rho \neq 0$ ,  $r=.45$ ,  
 $df=N-2=28-2=26$ ,  
 $r_{crit}=.374$

Reject  $H_0$  because  
 $r=.45 > r_{crit}=.374$



The SPSS p-value for  
 $r=.45$  with  $df=26$  is  
.0159. So reject  $H_0$   
because  $.0159 < \alpha=.05$ .

---

---

---

---

---

---

---

---

## Confidence Intervals for $\mu$

- Remember, interval estimation allows you to obtain an interval of potential values for a parameter
- For the problem about the ratio of estimated time to actual time for interrupted anagram solvers, we found  $\bar{X}=1.646$  for our sample mean
- We know that  $\bar{X}$  is a good (unbiased) estimate of  $\mu$  but we also know that  $\bar{X}$  has variability so it is unlikely that  $\mu=1.646$
- However, 1.646 should be close to  $\mu$
- Now we will see how to get an interval for  $\mu$  when we do not know  $\sigma^2$

---

---

---

---

---

---

---

## Confidence Intervals for $\mu$

- A confidence interval for  $\mu$  gives an interval of values around  $\bar{X}$  that are likely to include the true value of  $\mu$

- A 95% confidence interval for  $\mu$  is given by

$$\bar{X} - t_{\text{crit}}\left(\sqrt{\frac{s^2}{N}}\right) \text{ to } \bar{X} + t_{\text{crit}}\left(\sqrt{\frac{s^2}{N}}\right)$$

For the problem about the ratio of estimated time,  $\bar{X}=1.646$ ,  $N=10$ ,  $s^2=.3352$ ,  $df=N-1=9$ , and  $t_{\text{crit}}=\pm 2.262$

So the 95% confidence interval for  $\mu$  is

$$1.646 - 2.262\left(\sqrt{\frac{.3352}{10}}\right) \text{ to } 1.646 + 2.262\left(\sqrt{\frac{.3352}{10}}\right)$$

1.23 to 2.06

---

---

---

---

---

---

---